





# Full and Complete Binary Trees

Here are two important types of binary trees. Note that the definitions, while similar, are logically independent.

Definition: a binary tree  $T$  is *full* if each node is either a leaf or possesses exactly two child nodes.

Definition: a binary tree  $T$  with  $n$  levels is *complete* if all levels except possibly the last are completely full, and the last level has all its nodes to the left side.



Theorem: Let  $T$  be a nonempty, full binary tree Then:

- (a) If  $T$  has  $I$  internal nodes, the number of leaves is  $L = I + 1$ .
- (b) If  $T$  has  $I$  internal nodes, the total number of nodes is  $N = 2I + 1$ .
- (c) If  $T$  has a total of  $N$  nodes, the number of internal nodes is  $I = (N - 1)/2$ .
- (d) If  $T$  has a total of  $N$  nodes, the number of leaves is  $L = (N + 1)/2$ .
- (e) If  $T$  has  $L$  leaves, the total number of nodes is  $N = 2L - 1$ .
- (f) If  $T$  has  $L$  leaves, the number of internal nodes is  $I = L - 1$ .

Basically, this theorem says that the number of nodes  $N$ , the number of leaves  $L$ , and the number of internal nodes  $I$  are related in such a way that if you know any one of them, you can determine the other two.

proof of (a): We will use induction on the number of internal nodes,  $I$ . Let  $S$  be the set of all integers  $I \geq 0$  such that if  $T$  is a full binary tree with  $I$  internal nodes then  $T$  has  $I + 1$  leaf nodes.

For the base case, if  $I = 0$  then the tree must consist only of a root node, having no children because the tree is full. Hence there is 1 leaf node, and so  $0 \in S$ .

Now suppose that for some integer  $K \geq 0$ , every  $I$  from 0 through  $K$  is in  $S$ . That is, if  $T$  is a nonempty binary tree with  $I$  internal nodes, where  $0 \leq I \leq K$ , then  $T$  has  $I + 1$  leaf nodes.

Let  $T$  be a full binary tree with  $K + 1$  internal nodes. Then the root of  $T$  has two subtrees  $L$  and  $R$ ; suppose  $L$  and  $R$  have  $I_L$  and  $I_R$  internal nodes, respectively. Note that neither  $L$  nor  $R$  can be empty, and that every internal node in  $L$  and  $R$  must have been an internal node in  $T$ , and  $T$  had one additional internal node (the root), and so  $K + 1 = I_L + I_R + 1$ .

Now, by the induction hypothesis,  $L$  must have  $I_L + 1$  leaves and  $R$  must have  $I_R + 1$  leaves. Since every leaf in  $T$  must also be a leaf in either  $L$  or  $R$ ,  $T$  must have  $I_L + I_R + 2$  leaves.

Therefore, doing a tiny amount of algebra,  $T$  must have  $K + 2$  leaf nodes and so  $K + 1 \in S$ . Hence by Mathematical Induction,  $S = [0, \infty)$ .

QED

Theorem: Let  $T$  be a binary tree with  $\lambda$  levels. Then the number of leaves is at most  $2^{\lambda-1}$ .

proof: We will use strong induction on the number of levels,  $\lambda$ . Let  $S$  be the set of all integers  $\lambda \geq 1$  such that if  $T$  is a binary tree with  $\lambda$  levels then  $T$  has at most  $2^{\lambda-1}$  leaf nodes.

For the base case, if  $\lambda = 1$  then the tree must have one node (the root) and it must have no child nodes. Hence there is 1 leaf node (which is  $2^{\lambda-1}$  if  $\lambda = 1$ ), and so  $1 \in S$ .

Now suppose that for some integer  $K \geq 1$ , all the integers 1 through  $K$  are in  $S$ . That is, whenever a binary tree has  $M$  levels with  $M \leq K$ , it has at most  $2^{M-1}$  leaf nodes.

Let  $T$  be a binary tree with  $K + 1$  levels. If  $T$  has the maximum number of leaves,  $T$  consists of a root node and two nonempty subtrees, say  $S_1$  and  $S_2$ . Let  $S_1$  and  $S_2$  have  $M_1$  and  $M_2$  levels, respectively. Since  $M_1$  and  $M_2$  are between 1 and  $K$ , each is in  $S$  by the inductive assumption. Hence, the number of leaf nodes in  $S_1$  and  $S_2$  are at most  $2^{M_1-1}$  and  $2^{M_2-1}$ , respectively. Since all the leaves of  $T$  must be leaves of  $S_1$  or of  $S_2$ , the number of leaves in  $T$  is at most  $2^{M_1-1} + 2^{M_2-1}$  which is  $2^K$ . Therefore,  $K + 1$  is in  $S$ .

Hence by Mathematical Induction,  $S = [1, \infty)$ .

QED

Theorem: Let  $T$  be a binary tree. For every  $k \geq 0$ , there are no more than  $2^k$  nodes in level  $k$ .

Theorem: Let  $T$  be a binary tree with  $\lambda$  levels. Then  $T$  has no more than  $2^\lambda - 1$  nodes.

Theorem: Let  $T$  be a binary tree with  $N$  nodes. Then the number of levels is at least  $\lceil \log(N + 1) \rceil$ .

Theorem: Let  $T$  be a binary tree with  $L$  leaves. Then the number of levels is at least  $\lceil \log L \rceil + 1$ .